

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TA501318S

## HIBCH Mouse Monoclonal Antibody [Clone ID: OTI2A5]

### **Product data:**

| Product Type:           | Primary Antibodies   |
|-------------------------|--|
| Clone Name:             | OTI2A5   |
| Applications:           | FC, IF, WB   |
| Recommended Dilution:   | WB 1:2000, IF 1:100, FLOW 1:100  |
| Reactivity:             | Human  |
| Host:                   | Mouse  |
| lsotype:                | lgG1   |
| Clonality:              | Monoclonal   |
| Immunogen:              | Full length human recombinant protein of human HIBCH (NP_055177) produced in HEK293T<br>cell.  |
| Formulation:            | PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.   |
| Concentration:          | 0.5 mg/ml  |
| Purification:           | Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography<br>(protein A/G)   |
| Conjugation:            | Unconjugated   |
| Storage:                | Store at -20°C as received.  |
| Stability:              | Stable for 12 months from date of receipt.   |
| Predicted Protein Size: | 39.4 kDa   |
| Gene Name:              | 3-hydroxyisobutyryl-CoA hydrolase  |
| Database Link:          | <u>NP_055177</u><br><u>Entrez Gene 26275 Human</u><br><u>Q6NVY1</u>  |
| Background:             | This gene encodes the enzyme responsible for hydrolysis of both HIBYL-CoA and beta-<br>hydroxypropionyl-CoA. Mutations in this gene have been associated with 3-hyroxyisobutyryl-<br>CoA hydrolase deficiency. Alternative splicing results in multiple transcript variants. |
| Synonyms:               | HIBYLCOAH  |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

#### HIBCH Mouse Monoclonal Antibody [Clone ID: OTI2A5] - TA501318S


Protein Pathways: beta-Alanine metabolism, Metabolic pathways, Propanoate metabolism, Valine, leucine and isoleucine degradation

#### **Product images:**

158-106-79-

48-


35-23-



HepG2 HeLa HT29 A549 COS7 Jurkat MDCK PC12 MCF7


HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY HIBCH ([RC209814], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-HIBCH. Positive lysates [LY402324] (100ug) and [LC402324] (20ug) can be purchased separately from OriGene.

Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-HIBCH monoclonal antibody.



Anti-HIBCH mouse monoclonal antibody ([TA501318]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY HIBCH ([RC209814]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



HEK293T cells transfected with either [RC209814] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-HIBCH antibody ([TA501318]), and then analyzed by flow cytometry.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US