

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA501321BM

PANK2 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI3H9]

Product data:

Product Type:	Primary Antibodies
Clone Name:	ОТІЗН9
Applications:	FC, IF, IHC, WB
Recommended Dilution:	WB 1:1000~2000, IHC 1:50, IF 1:100, FLOW 1:100
Reactivity:	Human, Dog, Rat, Monkey, Mouse
Host:	Mouse
lsotype:	lgG2a
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human PANK2(NP_705902) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	HRP
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	59.1 kDa
Gene Name:	pantothenate kinase 2
Database Link:	<u>NP_705902</u> Entrez Gene 74450 MouseEntrez Gene 296167 RatEntrez Gene 477168 DogEntrez Gene 717639 MonkeyEntrez Gene 80025 Human <u>Q9BZ23</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


Serigene PANK2 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI3H9] – TA501321BM

- Background:This gene encodes a protein belonging to the pantothenate kinase family and is the only
member of that family to be expressed in mitochondria. Pantothenate kinase is a key
regulatory enzyme in the biosynthesis of coenzyme A (CoA) in bacteria and mammalian cells.
It catalyzes the first committed step in the universal biosynthetic pathway leading to CoA and
is itself subject to regulation through feedback inhibition by acyl CoA species. Mutations in
this gene are associated with HARP syndrome and pantothenate kinase-associated
neurodegeneration (PKAN), formerly Hallervorden-Spatz syndrome. Alternative splicing,
involving the use of alternate first exons, results in multiple transcripts encoding different
isoforms. [provided by RefSeq]
- Synonyms:C20orf48; HARP; HSS; NBIA1; PKANProtein Families:Druggable GenomeProtein Pathways:Metabolic pathways, Pantothenate and CoA biosynthesis


Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY PANK2 (Cat# [RC215676], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PANK2 (Cat# [TA501321]). Positive lysates [LY406998] (100ug) and [LC406998] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-PANK2 monoclonal antibody.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US 






Figure from citation: Western Blot of PANK2 protein level by using anti-PANK2 antibody in different mitochondrial compartments isolated from WT mouse brain and WT mouse fibroblasts. Dilution: 1:2000 <u>View Citation</u>




Immunohistochemical staining of paraffinembedded Human liver tissue within the normal limits using anti-PANK2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA501321], Dilution 1:50)

Immunohistochemical staining of paraffinembedded Human prostate tissue within the normal limits using anti-PANK2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA501321], Dilution 1:50)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US




Immunohistochemical staining of paraffinembedded Carcinoma of Human prostate tissue using anti-PANK2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA501321], Dilution 1:50)



Events PANK2 Anti-PANK2 mouse monoclonal antibody ([TA501321]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY PANK2 ([RC215676]).

HEK293T cells transfected with either [RC215676] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-PANK2 antibody ([TA501321]), and then analyzed by flow cytometry.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Flow cytometric Analysis of Jurkat cells, using anti-PANK2 antibody ([TA501321]), (Red), compared to a nonspecific negative control antibody (TA50011), (Blue).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US