

# Product datasheet for TA501326AM

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## SH3PX1 (SNX9) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI2F1]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI2F1

**Applications:** FC, IF, WB

**Recommended Dilution:** WB 1:1000~2000, IF 1:100, FLOW 1:100

Reactivity: Human, Monkey, Mouse, Rat

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human SNX9(NP\_057308) produced in HEK293T

cell

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 66.4 kDa

**Gene Name:** sorting nexin 9 **Database Link:** NP 057308

Entrez Gene 66616 MouseEntrez Gene 683687 RatEntrez Gene 706016 MonkeyEntrez Gene

51429 Human

Q9Y5X1



# SH3PX1 (SNX9) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI2F1] – TA501326AM


#### Background:

This gene encodes a member of the sorting nexin family. Members of this family contain a phox (PX) domain, which is a phosphoinositide binding domain, and are involved in intracellular trafficking. This protein does not contain a coiled coil region, like some family members, but does contain a SH3 domain near its N-terminus. This protein interacts with the cytoplasmic domains of the precursor but not the processed forms of a disintegrin and metalloprotease domain 9 and 15. This protein binds the beta-appendage domain of adaptor protein 2 and may function to assist adaptor protein 2 in its role at the plasma membrane. This protein interacts with activated Cdc42-associated kinase-2 to regulate the degradation of epidermal growth factor receptor protein. [provided by RefSeq]


Synonyms: SDP1; SH3PX1; SH3PXD3A; WISP

**Protein Families:** Druggable Genome

## **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY SNX9 ([RC202822], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-SNX9. Positive lysates [LY402520] (100ug) and [LC402520] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-SNX9 monoclonal antibody.






Western blot analysis of extracts (10ug) from 10 Human tissue by using anti-SNX9 monoclonal antibody at 1:200 (1: Testis; 2: Omentum; 3: Uterus; 4: Breast; 5: Brain; 6: Liver; 7: Ovary; 8: Thyroid gland; 9: colon;10: spleen).




Anti-SNX9 mouse monoclonal antibody ([TA501326]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY SNX9 ([RC202822] ).



HEK293T cells transfected with either [RC202822] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-SNX9 antibody ([TA501326]), and then analyzed by flow cytometry.





Flow cytometric Analysis of Jurkat cells, using anti-SNX9 antibody ([TA501326]), (Red), compared to a nonspecific negative control antibody (TA50011), (Blue).