

## **Product datasheet for TA501361AM**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## HIBCH Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI5C5]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI5C5

**Applications:** FC, IF, WB

Recommended Dilution: WB 1:2000, IF 1:100, FLOW 1:100

Reactivity: Human, Dog, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human HIBCH (NP\_055177) produced in HEK293T

cell.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 39.4 kDa

**Gene Name:** 3-hydroxyisobutyryl-CoA hydrolase

Database Link: NP 055177

Entrez Gene 301384 RatEntrez Gene 607040 DogEntrez Gene 26275 Human

Q6NVY1

Background: This gene encodes the enzyme responsible for hydrolysis of both HIBYL-CoA and beta-

hydroxypropionyl-CoA. Mutations in this gene have been associated with 3-hyroxyisobutyryl-

CoA hydrolase deficiency. Alternative splicing results in multiple transcript variants.

Synonyms: HIBYLCOAH







**Protein Pathways:** 

beta-Alanine metabolism, Metabolic pathways, Propanoate metabolism, Valine, leucine and isoleucine degradation


# **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY HIBCH ([RC209814], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-HIBCH. Positive lysates [LY402324] (100ug) and [LC402324] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-HIBCH monoclonal antibody.



Anti-HIBCH mouse monoclonal antibody ([TA501361]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY HIBCH ([RC209814]).



HEK293T cells transfected with either [RC209814] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-HIBCH antibody ([TA501361]), and then analyzed by flow cytometry.