

Product datasheet for TA501365AM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

TBXAS1 Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI1A12]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1A12
Applications: FC, IF, WB

Recommended Dilution: WB 1:2000, IF 1:100, FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human TBXAS1(NP_001052) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 60.5 kDa

Gene Name: thromboxane A synthase 1

Database Link: NP 001052

Entrez Gene 6916 Human

P24557





Background:


This gene encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. However, this protein is considered a member of the cytochrome P450 superfamily on the basis of sequence similarity rather than functional similarity. This endoplasmic reticulum membrane protein catalyzes the conversion of prostglandin H2 to thromboxane A2, a potent vasoconstrictor and inducer of platelet aggregation. The enzyme plays a role in several pathophysiological processes including hemostasis, cardiovascular disease, and stroke. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq]

Synonyms: BDPLT14; CYP5; CYP5A1; GHOSAL; THAS; TS; TXAS; TXS


Protein Families: Druggable Genome, P450

Protein Pathways: Arachidonic acid metabolism, Metabolic pathways


Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY TBXAS1 ([RC208028], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-TBXAS1. Positive lysates [LY420741] (100ug) and [LC420741] (20ug) can be purchased separately from OriGene.



Anti-TBXAS1 mouse monoclonal antibody ([TA501365]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY TBXAS1 ([RC208028]).





HEK293T cells transfected with either [RC208028] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-TBXAS1 antibody ([TA501365]), and then analyzed by flow cytometry.

Flow cytometric Analysis of Jurkat cells, using anti-TBXAS1 antibody ([TA501365]), (Red), compared to a nonspecific negative control antibody (TA50011), (Blue).