

## Product datasheet for TA501378AM

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## INDOL1 (IDO2) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI1A4]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: OTI1A4
Applications: FC, WB

Recommended Dilution: WB 1:500, FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human IDO2 (NP\_919270) produced in HEK293T

cell.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 46.9 kDa

**Gene Name:** indoleamine 2,3-dioxygenase 2

Database Link: NP 919270

Entrez Gene 169355 Human

Q6ZQW0

Background: Along with the enzymes encoded by the INDO (MIM 147435) and TDO2 (MIM 191070) genes,

the enzyme encoded by the INDOL1 gene metabolizes tryptophan in the kynurenine pathway

(Ball et al., 2007 [PubMed 17499941]). [supplied by OMIM]

Synonyms: INDOL1

**Protein Pathways:** Metabolic pathways, Tryptophan metabolism





## **Product images:**






Figure A, Western blot analysis of overexpressed lysates(15ug per lane) from HEK293T cells transfected with empty plasmid ([PS100001], NC), human IDO1 plasmid ([RC206592], hIDO1), human IDO2 plasmid ([RC223337], hIDO2) using anti-IDO2 antibody [TA501378](1:500). Figure B, Western blot analysis of the same samples as figure A with anti-DDK antibody ([TA180144], 1:1000)



HEK293T cells transfected with either [RC223337] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-IDO2 antibody ([TA501378]), and then analyzed by flow cytometry.



Flow cytometric Analysis of Jurkat cells, using anti-IDO2 antibody ([TA501378]), (Red), compared to a nonspecific negative control antibody, (Blue).