

# **Product datasheet for TA501473S**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## UCK (UCK1) Mouse Monoclonal Antibody [Clone ID: OTI4C2]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI4C2
Applications: FC, IF, WB

Recommended Dilution: WB 1:2000, IF 1:100, Flow: 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human UCK1 (NP\_113620) produced in HEK293T

cell

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 31.3 kDa

**Gene Name:** uridine-cytidine kinase 1

Database Link: NP 113620

Entrez Gene 83549 Human

Q9HA47

**Background:** Uridine/cytidine kinase-1 (EC 2.7.1.48) is a pyrimidine ribonucleoside kinase that catalyzes the

phosphorylation of uridine and cytidine to form uridine monophosphate (UMP) and cytidine monophosphate (CMP) (Van Rompay et al., 2001 [PubMed 11306702]). [supplied by OMIM]

Synonyms: URK1

**Protein Families:** Druggable Genome







**Protein Pathways:** 

Drug metabolism - other enzymes, Metabolic pathways, Pyrimidine metabolism


## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY UCK1 ([RC220876], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-UCK1. Positive lysates [LY403114] (100ug) and [LC403114] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-UCK1 monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human) (1:200).



Anti-UCK1 mouse monoclonal antibody ([TA501473]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY UCK1 ([RC220876]).





HEK293T cells transfected with either [RC220876] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-UCK1 antibody ([TA501473]), and then analyzed by flow cytometry.