

## **Product datasheet for TA501509**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Cytoplasmic dynein 1 light intermediate chain 1 (DYNC1LI1) Mouse Monoclonal Antibody [Clone ID: OTI2E8]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI2E8

**Applications:** FC, IF, WB

Recommended Dilution: WB 1:500, IF 1:100, FLOW 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human DYNC1LI1 (NP\_057225) produced in

HEK293T cell.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.34 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 56.4 kDa

**Gene Name:** dynein cytoplasmic 1 light intermediate chain 1

Database Link: NP 057225

Entrez Gene 235661 MouseEntrez Gene 252902 RatEntrez Gene 51143 Human

Q9Y6G9




# Cytoplasmic dynein 1 light intermediate chain 1 (DYNC1LI1) Mouse Monoclonal Antibody [Clone ID: OTI2E8] – TA501509

#### Background:


Acts as one of several non-catalytic accessory components of the cytoplasmic dynein 1 complex that are thought to be involved in linking dynein to cargos and to adapter proteins that regulate dynein function. Cytoplasmic dynein 1 acts as a motor for the intracellular retrograde motility of vesicles and organelles along microtubules. May play a role in binding dynein to membranous organelles or chromosomes. Probably involved in the microtubule-dependent transport of pericentrin. Is required for progress throuh the spindle assembly checkpoint. The phosphorylated form appears to be involved in the selective removal of MAD1L1 and MAD1L2 but not BUB1B from kinetochores.

Synonyms: DLC-A; DNCLI1; LIC1


### **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY DYNC1LI1 (Cat# [RC222010], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-DYNC1LI1(Cat# TA501509). Positive lysates [LY414168] (100ug) and [LC414168] (20ug) can be purchased separately from OriGene.




Anti-DYNC1LI1 mouse monoclonal antibody (TA501509) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY DYNC1LI1 ([RC222010]).



Flow cytometric Analysis of Jurkat cells, using anti-DYNC1LI1 antibody (TA501509), (Red), compared to a nonspecific negative control antibody (TA50011), (Blue).




Flow cytometric Analysis of Hela cells, using anti-DYNC1LI1 antibody (TA501509), (Red), compared to a nonspecific negative control antibody (TA50011), (Blue).



HEK293T cells transfected with either [RC222010] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-DYNC1LI1 antibody (TA501509), and then analyzed by flow cytometry.





Western blot analysis of extracts (10ug) from 2 different cell lines by using anti-DYNC1LI1 monoclonal antibody (1:200).