

Product datasheet for TA501543AM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

UCK (UCK1) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI5E9]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI5E9

Applications: FC, IF, WB

Recommended Dilution: WB 1:2000, IF 1:100, FLOW 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human UCK1 (NP_113620) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 31.3 kDa

Gene Name: uridine-cytidine kinase 1

Database Link: NP 113620

Entrez Gene 83549 Human

Q9HA47

Background: Uridine/cytidine kinase-1 (EC 2.7.1.48) is a pyrimidine ribonucleoside kinase that catalyzes the

phosphorylation of uridine and cytidine to form uridine monophosphate (UMP) and cytidine monophosphate (CMP) (Van Rompay et al., 2001 [PubMed 11306702]). [supplied by OMIM]

Synonyms: URK1

Protein Families: Druggable Genome







Protein Pathways:

Drug metabolism - other enzymes, Metabolic pathways, Pyrimidine metabolism


Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY UCK1 ([RC220876], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-UCK1. Positive lysates [LY403114] (100ug) and [LC403114] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (10ug) from 4 different cell lines by using anti-UCK1 monoclonal antibody (1:200).



Anti-UCK1 mouse monoclonal antibody ([TA501543]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY UCK1 ([RC220876]).



HEK293T cells transfected with either [RC220876] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-UCK1 antibody ([TA501543]), and then analyzed by flow cytometry.