

Product datasheet for **TA501543BM**

UCK (UCK1) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI5E9]

Product data:


Product Type:	Primary Antibodies
Clone Name:	OTI5E9
Applications:	FC, IF, WB
Recommended Dilution:	WB 1:2000, IF 1:100, FLOW 1:100
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human UCK1 (NP_113620) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	HRP
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	31.3 kDa
Gene Name:	uridine-cytidine kinase 1
Database Link:	NP_113620 Entrez Gene 83549 Human Q9HA47
Background:	Uridine/cytidine kinase-1 (EC 2.7.1.48) is a pyrimidine ribonucleoside kinase that catalyzes the phosphorylation of uridine and cytidine to form uridine monophosphate (UMP) and cytidine monophosphate (CMP) (Van Rompay et al., 2001 [PubMed 11306702]). [supplied by OMIM]
Synonyms:	URK1
Protein Families:	Druggable Genome




[View online »](#)

Protein Pathways: Drug metabolism - other enzymes, Metabolic pathways, Pyrimidine metabolism


Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY UCK1 ([RC220876], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-UCK1. Positive lysates [LY403114] (100ug) and [LC403114] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (10ug) from 4 different cell lines by using anti-UCK1 monoclonal antibody (1:200).



Anti-UCK1 mouse monoclonal antibody ([TA501543]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY UCK1 ([RC220876]).



HEK293T cells transfected with either [RC220876] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-UCK1 antibody ([TA501543]), and then analyzed by flow cytometry.