

## **Product datasheet for TA501544S**

## OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## IVD Mouse Monoclonal Antibody [Clone ID: OTI5A11]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: OTI5A11
Applications: FC, IF, WB

**Recommended Dilution:** WB 1:500~2000, IF 1:100, FLOW 1:100

Reactivity: Human, Dog, Rat, Monkey, Mouse

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human IVD (NP\_002612) produced in HEK293T

cell

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.66 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 46.5 kDa

**Gene Name:** isovaleryl-CoA dehydrogenase

Database Link: NP 002216

Entrez Gene 24513 RatEntrez Gene 56357 MouseEntrez Gene 100856316 DogEntrez Gene

702867 MonkeyEntrez Gene 3712 Human

P26440

**Background:** Isovaleryl-CoA dehydrogenase (IVD) is a mitochondrial matrix enzyme that catalyzes the third

step in leucine catabolism. The genetic deficiency of IVD results in an accumulation of isovaleric acid, which is toxic to the central nervous system and leads to isovaleric acidemia. Alternatively spliced transcript variants encoding different isoforms have been found for this

gene. [provided by RefSeq]






Synonyms: ACAD2; IVDH


**Protein Families:** Druggable Genome

**Protein Pathways:** Metabolic pathways, Valine, leucine and isoleucine degradation


## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY IVD (Cat# [RC229217], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-IVD(Cat# [TA501544]). Positive lysates [LY432239] (100ug) and [LC432239] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-IVD monoclonal antibody.



Anti-IVD mouse monoclonal antibody ([TA501544]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY IVD ([RC201077]).





HEK293T cells transfected with either [RC201077] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-IVD antibody ([TA501544]), and then analyzed by flow cytometry.