

Product datasheet for TA501546

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Ornithine Decarboxylase (ODC1) Mouse Monoclonal Antibody [Clone ID: OTI1G1]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1G1
Applications: FC, IF, WB

Recommended Dilution: WB 1:2000, IF 1:100, FLOW 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human ODC1 (NP_002530) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.54 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 51.0 kDa

Gene Name: ornithine decarboxylase 1

Database Link: NP 002530

Entrez Gene 18263 MouseEntrez Gene 24609 RatEntrez Gene 4953 Human

P11926

Background: This gene encodes the rate-limiting enzyme of the polyamine biosynthesis pathway which

catalyzes ornithine to putrescine. The activity level for the enzyme varies in response to growth-promoting stimuli and exhibits a high turnover rate in comparison to other

mammalian proteins. Originally localized to both chromosomes 2 and 7, the gene encoding this enzyme has been determined to be located on 2p25, with a pseudogene located on

7q31-qter. [provided by RefSeq]




Synonyms: BABS; NEDBA; NEDBIA; ODC


Protein Families: Druggable Genome

Protein Pathways: Arginine and proline metabolism, Glutathione metabolism, Metabolic pathways


Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY ODC1 ([RC206858], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ODC1. Positive lysates [LY400909] (100ug) and [LC400909] (20ug) can be purchased separately from OriGene.



Anti-ODC1 mouse monoclonal antibody (TA501546) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY ODC1 ([RC206858]).



HEK293T cells transfected with either [RC206858] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-ODC1 antibody (TA501546), and then analyzed by flow cytometry.