

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for TA501546AM

# Ornithine Decarboxylase (ODC1) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI1G1]

### **Product data:**


Product Type:	Primary Antibodies
Clone Name:	OTI1G1
Applications:	FC, IF, WB
Recommended Dilution:	WB 1:2000, IF 1:100, FLOW 1:100
Reactivity:	Human, Mouse, Rat
Host:	Mouse
lsotype:	lgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human ODC1 (NP_002530) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Biotin
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	51.0 kDa
Gene Name:	ornithine decarboxylase 1
Database Link:	<u>NP_002530</u> <u>Entrez Gene 18263 MouseEntrez Gene 24609 RatEntrez Gene 4953 Human</u> <u>P11926</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	Ornithine Decarboxylase (ODC1) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI1G1] – TA501546AM
Background:	This gene encodes the rate-limiting enzyme of the polyamine biosynthesis pathway which catalyzes ornithine to putrescine. The activity level for the enzyme varies in response to growth-promoting stimuli and exhibits a high turnover rate in comparison to other mammalian proteins. Originally localized to both chromosomes 2 and 7, the gene encoding this enzyme has been determined to be located on 2p25, with a pseudogene located on 7q31-qter. [provided by RefSeq]
Synonyms:	BABS; NEDBA; NEDBIA; ODC
Protein Families	: Druggable Genome
Protein Pathway	<i>ys:</i> Arginine and proline metabolism, Glutathione metabolism, Metabolic pathways

### **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY ODC1 ([RC206858], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ODC1. Positive lysates [LY400909] (100ug) and [LC400909] (20ug) can be purchased separately from OriGene.



Anti-ODC1 mouse monoclonal antibody ([TA501546]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY ODC1 ([RC206858]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US





HEK293T cells transfected with either [RC206858] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-ODC1 antibody ([TA501546]), and then analyzed by flow cytometry.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US