

# **Product datasheet for TA501546M**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Ornithine Decarboxylase (ODC1) Mouse Monoclonal Antibody [Clone ID: OTI1G1]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI1G1
Applications: FC, IF, WB

Recommended Dilution: WB 1:2000, IF 1:100, FLOW 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Full length human recombinant protein of human ODC1 (NP\_002530) produced in HEK293T

cell

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.54 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 51.0 kDa

**Gene Name:** ornithine decarboxylase 1

Database Link: NP 002530

Entrez Gene 18263 MouseEntrez Gene 24609 RatEntrez Gene 4953 Human

P11926

**Background:** This gene encodes the rate-limiting enzyme of the polyamine biosynthesis pathway which

catalyzes ornithine to putrescine. The activity level for the enzyme varies in response to growth-promoting stimuli and exhibits a high turnover rate in comparison to other

mammalian proteins. Originally localized to both chromosomes 2 and 7, the gene encoding

this enzyme has been determined to be located on 2p25, with a pseudogene located on

7q31-qter. [provided by RefSeq]






Synonyms: BABS; NEDBA; NEDBIA; ODC


**Protein Families:** Druggable Genome

**Protein Pathways:** Arginine and proline metabolism, Glutathione metabolism, Metabolic pathways


## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY ODC1 ([RC206858], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ODC1. Positive lysates [LY400909] (100ug) and [LC400909] (20ug) can be purchased separately from OriGene.



Anti-ODC1 mouse monoclonal antibody ([TA501546]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY ODC1 ([RC206858]).



HEK293T cells transfected with either [RC206858] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-ODC1 antibody ([TA501546]), and then analyzed by flow cytometry.