

## Product datasheet for **TA501578S**

### **B7-1 (CD80) Mouse Monoclonal Antibody [Clone ID: OTI1E9]**

#### **Product data:**

<b>Product Type:</b>	Primary Antibodies
<b>Clone Name:</b>	OTI1E9
<b>Applications:</b>	FC, IF, WB
<b>Recommended Dilution:</b>	WB 1:200~500, IF 1:100, FLOW 1:100
<b>Reactivity:</b>	Human, Dog, Rat, Monkey
<b>Host:</b>	Mouse
<b>Isotype:</b>	IgG1
<b>Clonality:</b>	Monoclonal
<b>Immunogen:</b>	Full length human recombinant protein of human CD80 (NP_005182) produced in HEK293T cell.
<b>Formulation:</b>	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
<b>Concentration:</b>	0.44 mg/ml
<b>Purification:</b>	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
<b>Conjugation:</b>	Unconjugated
<b>Storage:</b>	Store at -20°C as received.
<b>Stability:</b>	Stable for 12 months from date of receipt.
<b>Predicted Protein Size:</b>	29.3 kDa
<b>Gene Name:</b>	CD80 molecule
<b>Database Link:</b>	<a href="#">NP_005182</a> <a href="#">Entrez Gene 25408 Rat</a> <a href="#">Entrez Gene 403765 Dog</a> <a href="#">Entrez Gene 732518 Monkey</a> <a href="#">Entrez Gene 941 Human</a> <a href="#">P33681</a>
<b>Background:</b>	The B-lymphocyte activation antigen B7-1 (formerly referred to as B7) provides regulatory signals for T lymphocytes as a consequence of binding to the CD28 (MIM 186760) and CTLA4 (MIM 123890) ligands of T cells. [supplied by OMIM]
<b>Synonyms:</b>	B7; B7-1; B7.1; BB1; CD28LG; CD28LG1; LAB7




[View online »](#)


**Protein Families:** Druggable Genome, Transcription Factors, Transmembrane

**Protein Pathways:** Allograft rejection, Autoimmune thyroid disease, Cell adhesion molecules (CAMs), Graft-versus-host disease, Systemic lupus erythematosus, Toll-like receptor signaling pathway, Type I diabetes mellitus, Viral myocarditis


**Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY CD80 ([RC206540], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CD80.



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-CD80 monoclonal antibody.



Anti-CD80 mouse monoclonal antibody ([TA501578]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY CD80 ([RC206540]).



HEK293T cells transfected with either [RC206540] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-CD80 antibody ([TA501578]), and then analyzed by flow cytometry.