

Product datasheet for TA501669S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

B7-1 (CD80) Mouse Monoclonal Antibody [Clone ID: OTI7H3]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI7H3
Applications: FC, WB

Recommended Dilution: WB 1:2000

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human CD80 (NP_005182) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.64 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 29.3 kDa

Gene Name: CD80 molecule

Database Link: NP 005182

Entrez Gene 941 Human

P33681

Background: The B-lymphocyte activation antigen B7-1 (formerly referred to as B7) provides regulatory

signals for T lymphocytes as a consequence of binding to the CD28 (MIM 186760) and CTLA4

(MIM 123890) ligands of T cells. [supplied by OMIM]

Synonyms: B7; B7-1; B7.1; BB1; CD28LG; CD28LG1; LAB7

Protein Families: Druggable Genome, Transcription Factors, Transmembrane







Protein Pathways:

Allograft rejection, Autoimmune thyroid disease, Cell adhesion molecules (CAMs), Graft-versus-host disease, Systemic lupus erythematosus, Toll-like receptor signaling pathway, Type I diabetes mellitus, Viral myocarditis


Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY CD80 ([RC206540], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CD80.



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-CD80 monoclonal antibody.



HEK293T cells transfected with either [RC206540] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-CD80 antibody ([TA501669]), and then analyzed by flow cytometry (1:100).