

Product datasheet for TA501715

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

TUBA3E Mouse Monoclonal Antibody [Clone ID: OTI10B4]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI10B4
Applications: FC, WB

Recommended Dilution: WB 1:2000, FLOW 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human TUBA3E (NP_997195) produced in

HEK293T cell.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.56 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 49.7 kDa

Gene Name: tubulin alpha 3e

Database Link: NP 997195

Entrez Gene 112714 Human

Q6PEY2

Background: Tubulin is the major constituent of microtubules. It binds two moles of GTP, one at an

exchangeable site on the beta chain and one at a non-exchangeable site on the alpha-chain

(By similarity).

Synonyms: alpha 3e; tubulin


Protein Families: Druggable Genome






Protein Pathways: Gap junction, Pathogenic Escherichia coli infection

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY TUBA3E ([RC209279], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-TUBA3E. Positive lysates [LY404091] (100ug) and [LC404091] (20ug) can be purchased separately from OriGene.



HEK293T cells transfected with either [RC209279] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-TUBA3E antibody (TA501715), and then analyzed by flow cytometry.