

Product datasheet for TA501743

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

ARHGAP25 Mouse Monoclonal Antibody [Clone ID: OTI3H9]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI3H9

Applications: FC, IHC, WB

Recommended Dilution: WB 1:2000, IHC 1:150, FLOW 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human ARHGAP25 (NP_055697) produced in

HEK293T cell.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 72.3 kDa

Gene Name: Rho GTPase activating protein 25

Database Link: NP 055697

Entrez Gene 232201 MouseEntrez Gene 500246 RatEntrez Gene 9938 Human

P42331

Background: ARHGAPs, such as ARHGAP25, encode negative regulators of Rho GTPases (see ARHA; MIM

165390), which are implicated in actin remodeling, cell polarity, and cell migration (Katoh and


Katoh, 2004 [PubMed 15254788]). [supplied by OMIM]

Synonyms: HEL-S-308; KAIA0053







Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY ARHGAP25 ([RC217414], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ARHGAP25. Positive lysates [LY402385] (100ug) and [LC402385] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffinembedded Carcinoma of Human lung tissue using anti-ARHGAP25 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA501743, Dilution 1:50)




Immunohistochemical staining of paraffinembedded Human lymph node tissue within the normal limits using anti-ARHGAP25 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA501743, Dilution 1:50)





Immunohistochemical staining of paraffinembedded Human lymphoma tissue using anti-ARHGAP25 mouse monoclonal antibody. (Heatinduced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA501743, Dilution 1:50)



HEK293T cells transfected with either [RC217414] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-ARHGAP25 antibody (TA501743), and then analyzed by flow cytometry.