

# **Product datasheet for TA501762AM**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### PIK3AP1 Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI7G9]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI7G9
Applications: FC, IF, WB

**Recommended Dilution:** WB 1:500~2000, IF 1:100, FLOW 1:100

Reactivity: Human, Dog, Monkey, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human PIK3AP1 (NP\_689552) produced in

HEK293T cell.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

**Conjugation:** Biotin

Storage: Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 90.2 kDa

**Gene Name:** phosphoinositide-3-kinase adaptor protein 1

Database Link: NP 689522

Entrez Gene 294048 RatEntrez Gene 608653 DogEntrez Gene 705017 MonkeyEntrez Gene

<u>118788 Human</u>

Q6ZUJ8

**Background:** Involved in the activation of phosphoinositide 3-kinase (PI3K) in B-cells and in natural killer

(NK) cells. Couples B-cell antigen receptor (BCR) to PI3K activation by providing a docking site for the PI3K subunit PIK3R1, which contributes to B-cell development. Seems to have a complementary role with CD19 in PI3K activation (By similarity). May be involved in the

survival of mature B cells via activation of REL (By similarity).






Synonyms: BCAP


**Protein Families:** Druggable Genome

**Protein Pathways:** B cell receptor signaling pathway


## **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PIK3AP1 ([RC214125], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PIK3AP1. Positive lysates [LY403461] (100ug) and [LC403461] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-PIK3AP1 monoclonal antibody.



Anti-PIK3AP1 mouse monoclonal antibody ([TA501762]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY PIK3AP1 ([RC214125]).



HEK293T cells transfected with either [RC214125] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-PIK3AP1 antibody ([TA501762]), and then analyzed by flow cytometry.