

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA501864

Glutathione S Transferase theta 2 (GSTT2) Mouse Monoclonal Antibody [Clone ID: OTI7A1]

Product data:

Product Type:	Primary Antibodies		
Clone Name:	OTI7A1		
Applications:	FC, IF, WB		
Recommended Dilution:	WB 1:200~1000, IF 1:100, FLOW 1:100		
Reactivity:	Human		
Host:	Mouse		
lsotype:	lgG2a		
Clonality:	Monoclonal		
Immunogen:	Full length human recombinant protein of human GSTT2 (NP_000845) produced in HEK293T cell.		
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.		
Concentration:	0.58 mg/ml		
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)		
Conjugation:	Unconjugated		
Storage:	Store at -20°C as received.		
Stability:	Stable for 12 months from date of receipt.		
Predicted Protein Size:	27.3 kDa		
Gene Name:	glutathione S-transferase theta 2 (gene/pseudogene)		
Database Link:	<u>NP_000845</u> <u>Entrez Gene 2953 Human</u> <u>P0CG29</u>		




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	Glutathione S Transferase theta 2 (GSTT2) Mouse Monoclonal Antibody [Clone ID: OTI7A1] – TA501864
Background:	Glutathione S-transferase (GSTs) theta 2 (GSTT2) is a member of a superfamily of proteins that catalyze the conjugation of reduced glutathione to a variety of electrophilic and hydrophobic compounds. Human GSTs can be divided into five main classes: Alpha, Mu, Pi, Theta, and Zeta. The theta class members GSTT1 and GSTT2 share 55% amino acid sequence identity and both are thought to have an important role in human carcinogenesis. The theta genes have a similar structure, being composed of five exons with identical exon/intron boundaries. [provided by RefSeq]
Synonyms:	GSTT2B
Protein Pathway	vs: Drug metabolism - cytochrome P450, Glutathione metabolism, Metabolism of xenobiotics by cytochrome P450

Product images:

170	_	
130	-	
100	_	
70	_	
55	_	
40	_	
35	_	-
25	—	-
15	_	
10	_	

HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY GSTT2 ([RC200040], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-GSTT2. Positive lysates [LY424485] (100ug) and [LC424485] (20ug) can be purchased separately from OriGene.




Anti-GSTT2 mouse monoclonal antibody (TA501864) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY GSTT2 ([RC200040]).


This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Glutathione S Transferase theta 2 (GSTT2) Mouse Monoclonal Antibody [Clone ID: OTI7A1] – TA501864



Flow cytometric Analysis of Jurkat cells, using anti-GSTT2 antibody (TA501864), (Red), compared to a nonspecific negative control antibody (TA50011), (Blue).



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US