

Product datasheet for TA501864AM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Glutathione S Transferase theta 2 (GSTT2) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI7A1]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI7A1

Applications: FC, IF, WB

Recommended Dilution: WB 1:200~1000, IF 1:100, FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human GSTT2 (NP_000845) produced in HEK293T

cell.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 27.3 kDa

Gene Name: glutathione S-transferase theta 2 (gene/pseudogene)

Database Link: NP 000845

Entrez Gene 2953 Human

P0CG29





Glutathione S Transferase theta 2 (GSTT2) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI7A1] – TA501864AM

Background: Glutathione S-transferase (GSTs) theta 2 (GSTT2) is a member of a superfamily of proteins

that catalyze the conjugation of reduced glutathione to a variety of electrophilic and hydrophobic compounds. Human GSTs can be divided into five main classes: Alpha, Mu, Pi, Theta, and Zeta. The theta class members GSTT1 and GSTT2 share 55% amino acid sequence identity and both are thought to have an important role in human carcinogenesis. The theta genes have a similar structure, being composed of five exons with identical exon/intron


boundaries. [provided by RefSeq]

Synonyms: GSTT2B


Protein Pathways: Drug metabolism - cytochrome P450, Glutathione metabolism, Metabolism of xenobiotics by

cytochrome P450

Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY GSTT2 ([RC200040], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-GSTT2. Positive lysates [LY424485] (100ug) and [LC424485] (20ug) can be purchased separately from OriGene.




Anti-GSTT2 mouse monoclonal antibody ([TA501864]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY GSTT2 ([RC200040]).





Flow cytometric Analysis of Jurkat cells, using anti-GSTT2 antibody ([TA501864]), (Red), compared to a nonspecific negative control antibody (TA50011), (Blue).



Flow cytometric Analysis of Hela cells, using anti-GSTT2 antibody ([TA501864]), (Red), compared to a nonspecific negative control antibody (TA50011), (Blue).