

## **Product datasheet for TA501904**

#### 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

### ZIM2 Mouse Monoclonal Antibody [Clone ID: OTI5C7]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: OTI5C7
Applications: FC, WB

Recommended Dilution: WB 1:500, FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 1-150 and 428-527 of

human ZIM2 (NP\_056178) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.6 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 61 kDa

**Gene Name:** zinc finger imprinted 2

Database Link: NP 056178

Entrez Gene 23619 Human

O9NZV7




#### Background:

In human, ZIM2 and PEG3 (GeneID:5178) are two distinct genes that share a set of 5' exons and have a common promoter, and both genes are paternally expressed. Alternative splicing events connect the shared exons either with the remaining 4 exons unique to ZIM2, or with the remaining 2 exons unique to PEG3. This is in contrast to mouse and cow, where ZIM2 and PEG3 genes do not share exons in common, and the imprinting status of ZIM2 is also not conserved amongst mammals. Additional 5' alternatively spliced transcripts encoding the same protein have been found for the human ZIM2 gene. [provided by RefSeq]


Synonyms: ZNF656

**Protein Families:** Transcription Factors

# **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY ZIM2 ([RC211188], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ZIM2. Positive lysates [LY402427] (100ug) and [LC402427] (20ug) can be purchased separately from OriGene.



Flow cytometric Analysis of Jurkat cells, using anti-ZIM2 antibody (TA501904), (Red), compared to a nonspecific negative control antibody, (Blue).