

# Product datasheet for TA501959M

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### **HAO1 Mouse Monoclonal Antibody [Clone ID: OTI5A1]**

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI5A1

**Applications:** FC, IF, WB

**Recommended Dilution:** WB: 1:500-1:2000, IF: 1:100, Flow: 1:50-1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human HAO1 (NP\_060015) produced in HEK293T

cell

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.68 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

**Conjugation:** Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 40.7 kDa

**Gene Name:** hydroxyacid oxidase 1

Database Link: NP 060015

Entrez Gene 15112 MouseEntrez Gene 311446 RatEntrez Gene 54363 Human

Q9UJM8






#### Background:

This gene is one of three related genes that have 2-hydroxyacid oxidase activity yet differ in encoded protein amino acid sequence, tissue expression and substrate preference. Subcellular location of the encoded protein is the peroxisome. Specifically, this gene is expressed primarily in liver and pancreas and the encoded protein is most active on glycolate, a two-carbon substrate. The protein is also active on 2-hydroxy fatty acids. The transcript detected at high levels in pancreas may represent an alternatively spliced form or the use of a multiple near-consensus upstream polyadenylation site. [provided by RefSeq]. COMPLETENESS: full length.


**Synonyms:** GOX; GOX1; HAOX1

**Protein Pathways:** Glyoxylate and dicarboxylate metabolism, Metabolic pathways

# **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY HAO1 (Cat# [RC216834], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-HAO1(Cat# [TA501959]). Positive lysates [LY413692] (100ug) and [LC413692] (20ug) can be purchased separately from OriGene.



Anti-HAO1 mouse monoclonal antibody ([TA501959]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY HAO1 ([RC216834]).





HEK293T cells transfected with either [RC216834] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-HAO1 antibody ([TA501959]), and then analyzed by flow cytometry at 1:100