

## Product datasheet for **TA502055BM**

### **CALCOCO2 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI2B11]**

#### **Product data:**


<b>Product Type:</b>	Primary Antibodies
<b>Clone Name:</b>	OTI2B11
<b>Applications:</b>	FC, IF, WB
<b>Recommended Dilution:</b>	WB 1:2000, IF 1:100, FLOW 1:100
<b>Reactivity:</b>	Human
<b>Host:</b>	Mouse
<b>Isotype:</b>	IgG1
<b>Clonality:</b>	Monoclonal
<b>Immunogen:</b>	Full length human recombinant protein of human CALCOCO2 (NP_005822) produced in HEK293T cell.
<b>Formulation:</b>	PBS (pH 7.3) containing 1% BSA, 50% glycerol.
<b>Concentration:</b>	0.5 mg/ml
<b>Purification:</b>	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
<b>Conjugation:</b>	HRP
<b>Storage:</b>	Store at -20°C as received.
<b>Stability:</b>	Stable for 12 months from date of receipt.
<b>Predicted Protein Size:</b>	52.1 kDa
<b>Gene Name:</b>	calcium binding and coiled-coil domain 2
<b>Database Link:</b>	<a href="#">NP_005822</a> <a href="#">Entrez Gene 10241 Human</a> <a href="#">Q13137</a>
<b>Background:</b>	The protein encoded by this gene is a subunit of nuclear domain 10 (ND10) bodies. ND10 bodies are nuclear domains appearing immunohistochemically as ten dots per nucleus. They are believed to be associated with the nuclear matrix on the basis of their resistance to nuclease digestion and salt extraction. ND10 proteins are removed from the nucleus by herpes simplex virus-1 infection and may have a role in viral life cycles. [provided by RefSeq]. COMPLETENESS: complete on the 3' end.




[View online »](#)

Synonyms: NDP52


**Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY CALCOCO2 ([RC203843], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CALCOCO2. Positive lysates [LY417045] (100ug) and [LC417045] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (10ug) from 5 different cell lines by using anti-CALCOCO2 monoclonal antibody (1:200).



Anti-CALCOCO2 mouse monoclonal antibody ([TA502055]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY CALCOCO2 ([RC203843]).



HEK293T cells transfected with either [RC203843] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-CALCOCO2 antibody ([TA502055]), and then analyzed by flow cytometry.