

Product datasheet for TA502058BM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Annexin A3 (ANXA3) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI1F11]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1F11
Applications: IHC, WB

Recommended Dilution: WB 1:500~2000, IHC 1:150

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human ANXA3 (NP_005130) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 36.2 kDa

Gene Name: annexin A3

Database Link: NP 005130

Entrez Gene 11745 MouseEntrez Gene 25291 RatEntrez Gene 306 Human

P12429

Background: This gene encodes a member of the annexin family. Members of this calcium-dependent

phospholipid-binding protein family play a role in the regulation of cellular growth and in signal transduction pathways. This protein functions in the inhibition of phopholipase A2 and cleavage of inositol 1,2-cyclic phosphate to form inositol 1-phosphate. This protein may also

play a role in anti-coagulation. [provided by RefSeq, Jul 2008]







Synonyms: ANX3

Protein Families: Stem cell - Pluripotency


Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY ANXA3 (Cat# [RC201540], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ANXA3 (Cat# [TA502058]). Positive lysates [LY417500] (100ug) and [LC417500] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-ANXA3 monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human).



Western blot analysis of extracts (10ug) from 10 Human tissue by using anti-ANXA3 monoclonal antibody at 1:200 (1: Testis; 2: Omentum; 3: Uterus; 4: Breast; 5: Brain; 6: Liver; 7: Ovary; 8: Thyroid gland; 9: colon;10: spleen).






Figure from citation: Western blot analysis of ANXA3 protein level by using anti-ANXA3 antibody in CD133+ HCC cell lines. HCC cells, including HepG2, SMMC-7721, and Hep3B cells. Dilution: 1:1000 <u>View Citation</u>






Figure from citation: Western Blot of ANXA3 protein level by using anti-ANXA3 antibody in human HCC tissues and nontumorous tissues. Dilution: 1:1000 View Citation



Immunohistochemical staining of paraffinembedded Human lung tissue within the normal limits using anti-ANXA3 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502058])



Immunohistochemical staining of paraffinembedded Human prostate tissue within the normal limits using anti-ANXA3 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502058])






Figure from citation: Representative images of immunohistochemical staining of ANXA3 expression in human liver tissues: (C) negative ANXA3 expression in non-tumorous liver parenchyma and (D) weak, (E) moderate, and (F) strong ANXA3 immunoreactivity in HCC tissues. Dilution: 1:500 <u>View Citation</u>