

## **Product datasheet for TA502104S**

## **OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

## MYD88 Mouse Monoclonal Antibody [Clone ID: OTI1B4]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: OTI1B4

**Applications:** FC, IF, WB

Recommended Dilution: WB 1:2000, IF 1:100, FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human MYD88 (NP\_002459) produced in HEK293T

cell.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

**Conjugation:** Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 33.1 kDa

**Gene Name:** MYD88 innate immune signal transduction adaptor

Database Link: NP 002459

Entrez Gene 4615 Human

099836





Background:


This gene encodes a cytosolic adapter protein that plays a central role in the innate and adaptive immune response. This protein functions as an essential signal transducer in the interleukin-1 and Toll-like receptor signaling pathways. These pathways regulate that activation of numerous proinflammatory genes. The encoded protein consists of an N-terminal death domain and a C-terminal Toll-interleukin1 receptor domain. Patients with defects in this gene have an increased susceptibility to pyogenic bacterial infections. Alternate splicing results in multiple transcript variants. [provided by RefSeq]

Synonyms: MYD88D


**Protein Families:** Druggable Genome

**Protein Pathways:** Apoptosis, Toll-like receptor signaling pathway

## **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY MYD88 ([RC229151], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-MYD88. Positive lysates [LY432175] (100ug) and [LC432175] (20ug) can be purchased separately from OriGene.




Anti-MYD88 mouse monoclonal antibody ([TA502104]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY MYD88 ([RC202253]).






HEK293T cells transfected with either [RC202253] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-MYD88 antibody ([TA502104]), and then analyzed by flow cytometry.



Flow cytometric Analysis of Hela cells, using anti-MYD88 antibody ([TA502104]), (Red), compared to a nonspecific negative control antibody, (Blue).



Flow cytometric Analysis of Jurkat cells, using anti-MYD88 antibody ([TA502104]), (Red), compared to a nonspecific negative control antibody, (Blue).