

Product datasheet for TA502133M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

SQSTM1 Mouse Monoclonal Antibody [Clone ID: OTI4C8]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI4C8
Applications: FC, IF, WB

Recommended Dilution: WB 1:1000~2000, IF 1:100, FLOW 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human SQSTM1 (NP_003891) produced in

HEK293T cell.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.95 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 47.5 kDa

Gene Name: sequestosome 1

Database Link: NP 003891

Entrez Gene 18412 MouseEntrez Gene 113894 RatEntrez Gene 8878 Human

Q13501

Background: This gene encodes a multifunctional protein that binds ubiquitin and regulates activation of

the nuclear factor kappa-B (NF-kB) signaling pathway. The protein functions as a

scaffolding/adaptor protein in concert with TNF receptor-associated factor 6 to mediate activation of NF-kB in response to upstream signals. Alternatively spliced transcript variants encoding either the same or different isoforms have been identified for this gene. Mutations in this gene result in sporadic and familial Paget disease of bone. [provided by RefSeq]







Synonyms: A170; OSIL; p60; p62; p62B; PDB3; ZIP3

Protein Families: Druggable Genome, Transcription Factors


Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY SQSTM1 ([RC203214], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-SQSTM1. Positive lysates [LY401285] (100ug) and [LC401285] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-SQSTM1 monoclonal antibody.




Anti-SQSTM1 mouse monoclonal antibody ([TA502133]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY SQSTM1 ([RC203214]).





HEK293T cells transfected with either [RC203214] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-SQSTM1 antibody ([TA502133]), and then analyzed by flow cytometry.



Flow cytometric Analysis of Hela cells, using anti-SQSTM1 antibody ([TA502133]), (Red), compared to a nonspecific negative control antibody (TA50011), (Blue).