

Product datasheet for **TA502219M**

NMNAT1 Mouse Monoclonal Antibody [Clone ID: OTI1F7]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI1F7
Applications:	IF, WB
Recommended Dilution:	WB 1:2000, IF 1:100
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human NMNAT1 (NP_073624) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	31.8 kDa
Gene Name:	nicotinamide nucleotide adenyltransferase 1
Database Link:	NP_073624 Entrez Gene 64802 Human Q9HAN9
Background:	The coenzyme NAD and its derivatives are involved in hundreds of metabolic redox reactions and are utilized in protein ADP-ribosylation, histone deacetylation, and in some Ca(2+) signaling pathways. NMNAT (EC 2.7.7.1) is a central enzyme in NAD biosynthesis, catalyzing the condensation of nicotinamide mononucleotide (NMN) or nicotinic acid mononucleotide (NaMN) with the AMP moiety of ATP to form NAD or NaAD (Zhang et al., 2003 [PubMed 12574164]). [supplied by OMIM]




[View online »](#)


Synonyms: LCA9; NMNAT; PNAT1; SHILCA

Protein Pathways: Metabolic pathways, Nicotinate and nicotinamide metabolism


Product images:



Equivalent amounts of cell lysates (30 ug per lane) of wild-type HeLa cells (WT) and NMNAT1-Knockdown HeLa cells (KD) were separated by SDS-PAGE and immunoblotted with anti-NMNAT1 monoclonal antibody [TA502219] (1:2500). Then the blotted membrane was stripped and reprobed with anti-HSP90AA1 antibody as a loading control.



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY NMNAT1 ([RC204825], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NMNAT1 ([TA502219]). Positive lysates [LY402948] (100ug) and [LC402948] (20ug) can be purchased separately from OriGene.



Anti-NMNAT1 mouse monoclonal antibody ([TA502219]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY NMNAT1 ([RC204825]).