

Product datasheet for **TA502220BM**

NMNAT1 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI1G4]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI1G4
Applications:	FC, IHC, WB
Recommended Dilution:	WB 1:500~2000, IHC 1:150, FLOW 1:100
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human NMNAT1 (NP_073624) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	HRP
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	31.8 kDa
Gene Name:	nicotinamide nucleotide adenylyltransferase 1
Database Link:	NP_073624 Entrez Gene 64802 Human Q9HAN9
Background:	The coenzyme NAD and its derivatives are involved in hundreds of metabolic redox reactions and are utilized in protein ADP-ribosylation, histone deacetylation, and in some Ca(2+) signaling pathways. NMNAT (EC 2.7.7.1) is a central enzyme in NAD biosynthesis, catalyzing the condensation of nicotinamide mononucleotide (NMN) or nicotinic acid mononucleotide (NaMN) with the AMP moiety of ATP to form NAD or NaAD (Zhang et al., 2003 [PubMed 12574164]). [supplied by OMIM]




[View online »](#)


Synonyms: LCA9; NMNAT; PNAT1; SHILCA

Protein Pathways: Metabolic pathways, Nicotinate and nicotinamide metabolism


Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY NMNAT1 (Cat# [RC204825], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NMNAT1(Cat# [TA502220]). Positive lysates [LY402948] (100ug) and [LC402948] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-NMNAT1 monoclonal antibody.




Immunohistochemical staining of paraffin-embedded Human Kidney tissue within the normal limits using anti-NMNAT1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502220])




Immunohistochemical staining of paraffin-embedded Human Ovary tissue within the normal limits using anti-NMNAT1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502220])




Immunohistochemical staining of paraffin-embedded Human pancreas tissue within the normal limits using anti-NMNAT1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502220])




Immunohistochemical staining of paraffin-embedded Human endometrium tissue within the normal limits using anti-NMNAT1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502220])




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human endometrium tissue using anti-NMNAT1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502220])




Immunohistochemical staining of paraffin-embedded Human prostate tissue within the normal limits using anti-NMNAT1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502220])



Immunohistochemical staining of paraffin-embedded Carcinoma of Human prostate tissue using anti-NMNAT1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502220])



Immunohistochemical staining of paraffin-embedded Carcinoma of Human bladder tissue using anti-NMNAT1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502220])



HEK293T cells transfected with either [RC204825] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-NMNAT1 antibody ([TA502220]), and then analyzed by flow cytometry.