

# **Product datasheet for TA502232M**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## PPM1G Mouse Monoclonal Antibody [Clone ID: OTI1C12]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI1C12
Applications: IF, WB

**Reactivity:** WB 1:500~2000, IF 1:100 **Reactivity:** Human, Dog, Mouse, Rat

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human PPM1G (NP\_817092) produced in HEK293T

cell.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 59.1 kDa

**Gene Name:** protein phosphatase, Mg2+/Mn2+ dependent 1G

Database Link: NP 817092

Entrez Gene 475703 DogEntrez Gene 5496 Human

<u>O15355</u>

**Background:** The protein encoded by this gene is a member of the PP2C family of Ser/Thr protein

phosphatases. PP2C family members are known to be negative regulators of cell stress response pathways. This phosphatase is found to be responsible for the dephosphorylation of Pre-mRNA splicing factors, which is important for the formation of functional spliceosome. Studies of a similar gene in mice suggested a role of this phosphatase in regulating cell cycle

progression. [provided by RefSeq, Apr 2010]








**Synonyms:** PP2CG; PP2CGAMMA; PPP2CG

**Protein Families:** Druggable Genome, Phosphatase

#### **Product images:**







HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY PPM1G (Cat# [RC213685], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PPM1G(Cat# [TA502232]). Positive lysates [LY406071] (100ug) and [LC406071] (20ug) can be purchased separately from OriGene.

Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-PPM1G monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human).

Anti-PPM1G mouse monoclonal antibody ([TA502232]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY PPM1G ([RC213685]).