

Product datasheet for TA502264S

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

MMAB Mouse Monoclonal Antibody [Clone ID: OTI5G1]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI5G1

Applications: FC, IHC, WB

Recommended Dilution: WB 1:500, IHC 1:150, FLOW 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human MMAB (NP_443077) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.39 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 24.0 kDa

Gene Name: metabolism of cobalamin associated B

Database Link: NP 443077

Entrez Gene 326625 Human

O96EY8

Background: This gene encodes a protein that catalyzes the final step in the conversion of vitamin B(12)

into adenosylcobalamin (AdoCbl), a vitamin B12-containing coenzyme for methylmalonyl-CoA

mutase. Mutations in the gene are the cause of vitamin B12-dependent methylmalonic

aciduria linked to the cblB complementation group. [provided by RefSeq]

Synonyms: ATR; cblB; CFAP23; cob







Protein Pathways:

Metabolic pathways, Porphyrin and chlorophyll metabolism


Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY MMAB ([RC204290], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-MMAB. Positive lysates [LY409455] (100ug) and [LC409455] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffinembedded Human liver tissue within the normal limits using anti-MMAB mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502264])




Immunohistochemical staining of paraffinembedded Carcinoma of Human lung tissue using anti-MMAB mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502264])





Immunohistochemical staining of paraffinembedded Human Ovary tissue within the normal limits using anti-MMAB mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502264])



HEK293T cells transfected with either [RC204290] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-MMAB antibody ([TA502264]), and then analyzed by flow cytometry.