

## Product datasheet for TA502278M

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### MMAA Mouse Monoclonal Antibody [Clone ID: OTI3A3]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI3A3
Applications: IHC, WB

Recommended Dilution: WB 1:200~500, IHC 1:150

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Full length human recombinant protein of human MMAA (NP\_785454) produced in HEK293T

cell

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.75 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 39.0 kDa

**Gene Name:** metabolism of cobalamin associated A

Database Link: NP 758454

Entrez Gene 166785 Human

**Q8IVH4** 

**Background:** The protein encoded by this gene is involved in the translocation of cobalamin into the

mitochondrion, where it is used in the final steps of adenosylcobalamin synthesis.

Adenosylcobalamin is a coenzyme required for the activity of methylmalonyl-CoA mutase.

Defects in this gene are a cause of methylmalonic aciduria. [provided by RefSeq]


Synonyms: cblA






**Protein Families:** Druggable Genome


# **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY MMAA ([RC223205], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-MMAA. Positive lysates [LY403540] (100ug) and [LC403540] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-MMAA monoclonal antibody.



Immunohistochemical staining of paraffinembedded Human Kidney tissue within the normal limits using anti-MMAA mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.