

Product datasheet for TA502297M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

ACY3 Mouse Monoclonal Antibody [Clone ID: OTI2G2]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2G2

Applications: FC, IHC, WB

Recommended Dilution: WB 1:200~500, IHC 1:150, FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human ACY3 (NP_542389) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.56 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 35.1 kDa

Gene Name: aminoacylase 3

Database Link: NP 542389

Entrez Gene 91703 Human


Q96HD9

Synonyms: ACY-3; HCBP1


Protein Pathways: Alanine, aspartate and glutamate metabolism, Histidine metabolism




Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY ACY3 ([RC202287], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ACY3. Positive lysates [LY408962] (100ug) and [LC408962] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-ACY3 monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human).



Immunohistochemical staining of paraffinembedded Human Kidney tissue within the normal limits using anti-ACY3 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



HEK293T cells transfected with either [RC202287] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-ACY3 antibody ([TA502297]), and then analyzed by flow cytometry.