

Product datasheet for TA502390S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

VCAM1 Mouse Monoclonal Antibody [Clone ID: OTI4A10]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI4A10
Applications: FC, WB

Recommended Dilution: WB 1:200~500, FLOW 1:100

Reactivity: Human, Dog, Monkey

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human VCAM1 (NP_001069) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.58 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 78.7 kDa

Gene Name: vascular cell adhesion molecule 1

Database Link: NP 001069

Entrez Gene 403982 DogEntrez Gene 712036 MonkeyEntrez Gene 7412 Human

P19320

Background: This gene is a member of the Ig superfamily and encodes a cell surface sialoglycoprotein

expressed by cytokine-activated endothelium. This type I membrane protein mediates leukocyte-endothelial cell adhesion and signal transduction, and may play a role in the development of artherosclerosis and rheumatoid arthritis. Two alternatively spliced transcripts encoding different isoforms have been described for this gene. [provided by

RefSeq1






Synonyms: CD106; INCAM-100


Protein Families: Druggable Genome, ES Cell Differentiation/IPS, Transmembrane

Protein Pathways: Cell adhesion molecules (CAMs), Leukocyte transendothelial migration


Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY VCAM1 ([RC209761], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-VCAM1. Positive lysates [LY421491] (100ug) and [LC421491] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-VCAM1 monoclonal antibody.



Flow cytometric Analysis of Hela cells, using anti-VCAM1 antibody ([TA502390]), (Red), compared to a nonspecific negative control antibody (TA50011), (Blue).