

## **Product datasheet for TA502455**

### **OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

# POLR2J2 Mouse Monoclonal Antibody [Clone ID: OTI6E9]

### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI6E9
Applications: FC, WB

Recommended Dilution: WB 1:2000, FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human POLR2J2 (NP\_116581) produced in

HEK293T cell.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.94 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 12.9 kDa

**Gene Name:** RNA polymerase II subunit J2

Database Link: NP 116581

Entrez Gene 246721 Human

O9GZM3





#### Background:

This gene is a member of the RNA polymerase II subunit 11 gene family, which includes three genes in a cluster on chromosome 7q22.1 and a pseudogene on chromosome 7p13. The founding member of this family, DNA directed RNA polymerase II polypeptide J, has been shown to encode a subunit of RNA polymerase II, the polymerase responsible for synthesizing messenger RNA in eukaryotes. This locus produces multiple, alternatively spliced transcripts that potentially express isoforms with distinct C-termini compared to DNA directed RNA polymerase II polypeptide J. Most or all variants are spliced to include additional non-coding exons at the 3' end which makes them candidates for nonsense-mediated decay (NMD). Consequently, it is not known if this locus expresses a protein or proteins in vivo. [provided by RefSeq]


Synonyms: HRPB11B; RPB11b1

Protein Families: Transcription Factors


Protein Pathways: Huntington's disease, Metabolic pathways, Purine metabolism, Pyrimidine metabolism, RNA

polymerase

### **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY POLR2J2 ([RC224755], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-POLR2J2. Positive lysates [LY409843] (100ug) and [LC409843] (20ug) can be purchased separately from OriGene.



HEK293T cells transfected with either [RC224755] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-POLR2J2 antibody (TA502455), and then analyzed by flow cytometry.