

Product datasheet for TA502467S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PIH1D2 Mouse Monoclonal Antibody [Clone ID: OTI4A10]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI4A10

Applications: FC, IHC, WB

Recommended Dilution: WB 1:1000, IHC 1:150, FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 1-315 of human

PIH1D2(NP_620144) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.83 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 35.8 kDa

Gene Name: PIH1 domain containing 2

Database Link: NP 620144

Entrez Gene 120379 Human


Q8WWB5

Synonyms: PIH1 domain containing 2







Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY PIH1D2 (Cat# [RC205039], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PIH1D2(Cat# [TA502467]). Positive lysates [LY408526] (100ug) and [LC408526] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffinembedded Human endometrium tissue within the normal limits using anti-PIH1D2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502467])




Immunohistochemical staining of paraffinembedded Carcinoma of Human bladder tissue using anti-PIH1D2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502467])






Immunohistochemical staining of paraffinembedded Human tonsil within the normal limits using anti-PIH1D2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502467])



Flow cytometric Analysis of Hela cells, using anti-PIH1D2 antibody ([TA502467]), (Red), compared to a nonspecific negative control antibody, (Blue).



Flow cytometric Analysis of Jurkat cells, using anti-PIH1D2 antibody ([TA502467]), (Red), compared to a nonspecific negative control antibody, (Blue).