

## Product datasheet for **TA502482BM**

### NDUFB9 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI3A1]


#### Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI3A1
Applications:	IHC, WB
Recommended Dilution:	WB 1:2000, IHC 1:150
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Human recombinant protein fragment corresponding to amino acids 3-179 of human NDUFB9(NP_004996) produced in E.coli.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	HRP
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	21.7 kDa
Gene Name:	NADH:ubiquinone oxidoreductase subunit B9
Database Link:	<a href="#">NP_004996</a> <a href="#">Entrez Gene 66218 Mouse</a> <a href="#">Entrez Gene 299954 Rat</a> <a href="#">Entrez Gene 4715 Human</a> <a href="#">Q9Y6M9</a>
Synonyms:	B22; CI-B22; LYRM3; UQOR22
Protein Pathways:	Alzheimer's disease, Huntington's disease, Metabolic pathways, Oxidative phosphorylation, Parkinson's disease




[View online »](#)


**Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY NDUFB9 ([RC200223], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NDUFB9. Positive lysates [LY417578] (100ug) and [LC417578] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human colon tissue using anti-NDUFB9 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502482])




Immunohistochemical staining of paraffin-embedded Human Kidney tissue within the normal limits using anti-NDUFB9 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502482])




Immunohistochemical staining of paraffin-embedded Human liver tissue within the normal limits using anti-NDUFB9 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502482])




Immunohistochemical staining of paraffin-embedded Carcinoma of Human liver tissue using anti-NDUFB9 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502482])




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human ovary tissue using anti-NDUFB9 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502482])



Immunohistochemical staining of paraffin-embedded Human pancreas tissue within the normal limits using anti-NDUFB9 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502482])



Immunohistochemical staining of paraffin-embedded Carcinoma of Human thyroid tissue using anti-NDUFB9 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502482])



Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human endometrium tissue using anti-NDUFB9 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502482])