

Product datasheet for **TA502492BM**

VASP Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI1C3]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI1C3
Applications:	FC, IF, WB
Recommended Dilution:	WB 1:500~2000, IF 1:100, FLOW 1:100
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human VASP (NP_003361) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	HRP
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	39.6 kDa
Gene Name:	vasodilator stimulated phosphoprotein
Database Link:	NP_003361 Entrez Gene 22323 Mouse Entrez Gene 361517 Rat Entrez Gene 7408 Human P50552



[View online »](#)

Background:

Vasodilator-stimulated phosphoprotein (VASP) is a member of the Ena-VASP protein family. Ena-VASP family members contain an EHV1 N-terminal domain that binds proteins containing E/DFPPPPXD/E motifs and targets Ena-VASP proteins to focal adhesions. In the mid-region of the protein, family members have a proline-rich domain that binds SH3 and WW domain-containing proteins. Their C-terminal EVH2 domain mediates tetramerization and binds both G and F actin. VASP is associated with filamentous actin formation and likely plays a widespread role in cell adhesion and motility. VASP may also be involved in the intracellular signaling pathways that regulate integrin-extracellular matrix interactions. VASP is regulated by the cyclic nucleotide-dependent kinases PKA and PKG. [provided by RefSeq]

Synonyms:


vasodilator-stimulated phosphoprotein

Protein Families:


Druggable Genome, Stem cell - Pluripotency

Protein Pathways:


Fc gamma R-mediated phagocytosis, Focal adhesion, Leukocyte transendothelial migration

Product images:


HEK293T cells transfected with either [RC203544] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-VASP antibody ([TA502492]), and then analyzed by flow cytometry.




Anti-VASP mouse monoclonal antibody ([TA502492]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY VASP ([RC203544]).



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY VASP ([RC203544], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-VASP. Positive lysates [LY401150] (100ug) and [LC401150] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-VASP monoclonal antibody.



Equivalent amounts of cell lysates (10 ug per lane) of wild-type HeLa cells (WT, Cat# LC810HELA) and VASP-Knockout HeLa cells (KO, Cat# [LC832038]) were separated by SDS-PAGE and immunoblotted with anti-VASP monoclonal antibody [TA502492] (1:500). Then the blotted membrane was stripped and reprobed with anti-HSP90 antibody as a loading control.