

## Product datasheet for **TA502624BM**


### NNMT Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI3D8]

#### Product data:


Product Type:	Primary Antibodies
Clone Name:	OTI3D8
Applications:	FC, IF, IHC, WB
Recommended Dilution:	WB 1:500~2000, IHC 1:150, IF 1:100, FLOW 1:100
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG2b
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human NNMT(NP_006160) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	HRP
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	29.4 kDa
Gene Name:	nicotinamide N-methyltransferase
Database Link:	<a href="#">NP_006160</a> <a href="#">Entrez Gene 18113 Mouse</a> <a href="#">Entrez Gene 300691 Rat</a> <a href="#">Entrez Gene 4837 Human</a> <a href="#">P40261</a>
Background:	N-methylation is one method by which drug and other xenobiotic compounds are metabolized by the liver. This gene encodes the protein responsible for this enzymatic activity which uses S-adenosyl methionine as the methyl donor. [provided by RefSeq]
Synonyms:	nicotinamide N-methyltransferase
Protein Pathways:	Metabolic pathways, Nicotinate and nicotinamide metabolism


[View online »](#)


## Product images:




Western blot analysis of extracts (10ug) from 9 Human tissue by using anti-NNMT monoclonal antibody at 1:200 (1: Testis; 2: Omentum; 3: Uterus; 4: Breast; 5: Brain; 6: Liver; 7: Ovary; 8: Thyroid gland; 9: colon).




Equivalent amounts of cell lysates (10 ug per lane) of wild-type HeLa cells (WT, Cat# LC810HELA) and NNMT-Knockout HeLa cells were separated by SDS-PAGE and immunoblotted with anti-NNMT monoclonal antibody [TA502624] (1:500). Then the blotted membrane was stripped and reprobed with anti-HSP90 antibody as a loading control.




Western blot analysis of overexpressed lysates(15ug per lane) from HEK293T cells transfected with empty plasmid ([PS100001], lane 1) , human NNMT plasmid ([RC200641], lane 2), mouse NNMT plasmid ([MR203499], lane 3) using anti-NNMT antibody [TA502624] (1:500).




Western blot analysis of extracts (50ug per lane) from 3 cell lines lysates by using anti-NNMT monoclonal antibody([TA502624], 1:500)




Immunohistochemical staining of paraffin-embedded Human liver tissue within the normal limits using anti-NNMT mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502624])



Immunohistochemical staining of paraffin-embedded Human pancreas tissue within the normal limits using anti-NNMT mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502624])




Immunohistochemical staining of paraffin-embedded Carcinoma of Human thyroid tissue using anti-NNMT mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502624])




Immunohistochemical staining of paraffin-embedded Human endometrium tissue within the normal limits using anti-NNMT mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502624])







Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human endometrium tissue using anti-NNMT mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502624])



Immunohistochemical staining of paraffin-embedded Human prostate tissue within the normal limits using anti-NNMT mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502624])



Anti-NNMT mouse monoclonal antibody ([TA502624]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY NNMT ([RC200641]).



Flow cytometric Analysis of Jurkat cells, using anti-NNMT antibody ([TA502624]), (Red), compared to a nonspecific negative control antibody, (Blue).