

Product datasheet for TA502641BM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Destrin (DSTN) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI1C8]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1C8

Applications: FC, IF, IHC, WB

Recommended Dilution: WB 1:500~2000, IHC 1:150, IF 1:100, FLOW 1:100

Reactivity: Human, Mouse

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human DSTN (NP_006861) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 18.3 kDa

Gene Name: destrin, actin depolymerizing factor

Database Link: NP 006861

Entrez Gene 11034 Human

P60981

Background: The product of this gene belongs to the actin-binding proteins ADF family. This family of

proteins is responsible for enhancing the turnover rate of actin in vivo. This gene encodes the

actin depolymerizing protein that severs actin filaments (F-actin) and binds to actin

monomers (G-actin). Two transcript variants encoding distinct isoforms have been identified


for this gene. [provided by RefSeq]






Synonyms: ACTDP; ADF; bA462D18.2; HEL32


Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY DSTN ([RC203419], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-DSTN. Positive lysates [LY402051] (100ug) and [LC402051] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-DSTN monoclonal antibody.




Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-DSTN mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502641])






Immunohistochemical staining of paraffinembedded Carcinoma of Human thyroid tissue using anti-DSTN mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502641])




Immunohistochemical staining of paraffinembedded Human prostate tissue within the normal limits using anti-DSTN mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502641])




Immunohistochemical staining of paraffinembedded Carcinoma of Human bladder tissue using anti-DSTN mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502641])






Anti-DSTN mouse monoclonal antibody ([TA502641]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY DSTN ([RC203419]).




HEK293T cells transfected with either [RC203419] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-DSTN antibody ([TA502641]), and then analyzed by flow cytometry.



Flow cytometric Analysis of Hela cells, using anti-DSTN antibody ([TA502641]), (Red), compared to a nonspecific negative control antibody (TA50011), (Blue).





Flow cytometric Analysis of Jurkat cells, using anti-DSTN antibody ([TA502641]), (Red), compared to a nonspecific negative control antibody (TA50011), (Blue).