

## Product datasheet for TA502662M

## 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc.

## KCTD14 Mouse Monoclonal Antibody [Clone ID: OTI2B9]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: OTI2B9

**Applications:** FC, IF, IHC, WB

**Recommended Dilution:** WB 1:2000, IHC 1:150, IF 1:100, FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human KCTD14 (NP\_076419) produced in

HEK293T cell.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.77 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 29.4 kDa

**Gene Name:** potassium channel tetramerization domain containing 14

Database Link: NP 076419

Entrez Gene 65987 Human

Q9BQ13


Synonyms: MGC2376

**Protein Families:** Ion Channels: Other






## **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY KCTD14 ([RC200745], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-KCTD14. Positive lysates [LY411435] (100ug) and [LC411435] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-KCTD14 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.






Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human endometrium tissue using anti-KCTD14 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.




Anti-KCTD14 mouse monoclonal antibody ([TA502662]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY KCTD14 ([RC200745]).



Flow cytometric Analysis of Hela cells, using anti-KCTD14 antibody ([TA502662]), (Red), compared to a nonspecific negative control antibody (TA50011), (Blue).





Flow cytometric Analysis of Jurkat cells, using anti-KCTD14 antibody ([TA502662]), (Red), compared to a nonspecific negative control antibody (TA50011), (Blue).