

## Product datasheet for **TA502717BM**


### **RACK1 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI2D8]**

#### **Product data:**


Product Type:	Primary Antibodies
Clone Name:	OTI2D8
Applications:	FC, IHC, WB
Recommended Dilution:	WB 1:500, IHC 1:150, FLOW 1:100
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human GNB2L1 (NP_006089) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	HRP
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	34.9 kDa
Gene Name:	receptor for activated C kinase 1
Database Link:	<a href="#">NP_006089</a> <a href="#">Entrez Gene 10399 Human P63244</a>
Synonyms:	Gnb2-rs1; GNB2L1; H12.3; HLC-7; PIG21
Protein Families:	Druggable Genome




[View online »](#)

**Product images:**


HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY GNB2L1 (Cat# [RC205092], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-GNB2L1 (Cat# [TA502717]). Positive lysates [LY401838] (100ug) and [LC401838] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human breast tissue using anti-GNB2L1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502717])




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human colon tissue using anti-GNB2L1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502717])




Immunohistochemical staining of paraffin-embedded Human Kidney tissue within the normal limits using anti-GNB2L1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502717])




Immunohistochemical staining of paraffin-embedded Carcinoma of Human prostate tissue using anti-GNB2L1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502717])




Immunohistochemical staining of paraffin-embedded Carcinoma of Human bladder tissue using anti-GNB2L1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502717])




Immunohistochemical staining of paraffin-embedded Human lymph node tissue within the normal limits using anti-GNB2L1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502717])



HEK293T cells transfected with either [RC205092] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-GNB2L1 antibody ([TA502717]), and then analyzed by flow cytometry.



Flow cytometric Analysis of HeLa cells, using anti-GNB2L1 antibody ([TA502717]), (Red), compared to a nonspecific negative control antibody, (Blue).



Flow cytometric Analysis of Jurkat cells, using anti-GNB2L1 antibody ([TA502717]), (Red), compared to a nonspecific negative control antibody, (Blue).