

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for TA502728AM

## SSX1 Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI2D12]

## **Product data:**

Product Type:	Primary Antibodies
Clone Name:	OTI2D12
Applications:	FC, IF, WB
Recommended Dilution:	WB 1:2000, IF 1:100, FLOW 1:100
Reactivity:	Human
Host:	Mouse
lsotype:	lgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human SSX1 (NP_005626) produced in E.coli.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Biotin
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	21.8 kDa
Gene Name:	SSX family member 1
Database Link:	<u>NP_005626</u> <u>Entrez Gene 6756 Human</u> <u>Q16384</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

#### SSX1 Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI2D12] – TA502728AM


Background:The product of this gene belongs to the family of highly homologous synovial sarcoma X (SSX)<br/>breakpoint proteins. These proteins may function as transcriptional repressors. They are also<br/>capable of eliciting spontaneously humoral and cellular immune responses in cancer<br/>patients, and are potentially useful targets in cancer vaccine-based immunotherapy. SSX1,<br/>SSX2 and SSX4 genes have been involved in the t(X;18) translocation characteristically found<br/>in all synovial sarcomas. This translocation results in the fusion of the synovial sarcoma<br/>translocation gene on chromosome 18 to one of the SSX genes on chromosome X. The<br/>encoded hybrid proteins are probably responsible for transforming activity. [provided by<br/>RefSeq]

Synonyms:	CT5.1; SSRC


Protein Families:

# Transcription Factors

### **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY SSX1 ([RC201600], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-SSX1. Positive lysates [LY417166] (100ug) and [LC417166] (20ug) can be purchased separately from OriGene.



Anti-SSX1 mouse monoclonal antibody ([TA502728]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY SSX1 ([RC201600]).


This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



SSX1

HEK293T cells transfected with either [RC201600] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-SSX1 antibody ([TA502728]), and then analyzed by flow cytometry.

Flow cytometric Analysis of Hela cells, using anti-SSX1 antibody ([TA502728]), (Red), compared to a nonspecific negative control antibody (TA50011), (Blue).



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US