

Product datasheet for TA502804M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

ALDOB Mouse Monoclonal Antibody [Clone ID: OTI2A3]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2A3

Applications: Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: lgG1

Clonality: Monoclonal

Full length human recombinant protein of human ALDOB (NP_000026) produced in HEK293T Immunogen:

WB

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.42 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Store at -20°C as received. Storage:

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 39.3 kDa

Gene Name: aldolase, fructose-bisphosphate B

Database Link: NP 000026

Entrez Gene 24190 RatEntrez Gene 230163 MouseEntrez Gene 229 Human

P05062





Background:

Fructose-1,6-bisphosphate aldolase (EC 4.1.2.13) is a tetrameric glycolytic enzyme that catalyzes the reversible conversion of fructose-1,6-bisphosphate to glyceraldehyde 3-phosphate and dihydroxyacetone phosphate. Vertebrates have 3 aldolase isozymes which are distinguished by their electrophoretic and catalytic properties. Differences indicate that aldolases A, B, and C are distinct proteins, the products of a family of related 'housekeeping' genes exhibiting developmentally regulated expression of the different isozymes. The developing embryo produces aldolase A, which is produced in even greater amounts in adult muscle where it can be as much as 5% of total cellular protein. In adult liver, kidney and intestine, aldolase A expression is repressed and aldolase B is produced. In brain and other nervous tissue, aldolase A and C are expressed about equally. There is a high degree of homology between aldolase A and C. Defects in ALDOB cause hereditary fructose intolerance. [provided by RefSeq]


Synonyms: ALDB; ALDO2

Protein Families: Druggable Genome

Protein Pathways: Fructose and mannose metabolism, Glycolysis / Gluconeogenesis, Metabolic pathways,


Pentose phosphate pathway

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY ALDOB ([RC220062], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ALDOB. Positive lysates [LY400007] (100ug) and [LC400007] (20ug) can be purchased separately from OriGene.





Western blot analysis of extracts (10ug) from 4 different cell lines by using anti-ALDOB monoclonal antibody (1:200).