

Product datasheet for TA502841BM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

ALDH1A3 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI4E8]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI4E8
Applications: FC, WB

Recommended Dilution: WB 1:200~2000, FLOW 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 1-100 and 413-512 of

human ALDH1A3 (NP_000648) produced in E.coli.

Formulation: PBS (pH 7.4) containing 1% BSA, 50% glycerol.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 55.9 kDa

Gene Name: aldehyde dehydrogenase 1 family member A3

Database Link: NP 000684

Entrez Gene 220 Human

P47895

Background: Aldehyde dehydrogenase isozymes are thought to play a major role in the detoxification of

aldehydes generated by alcohol metabolism and lipid peroxidation. The enzyme encoded by this gene uses retinal as a substrate, either in a free or cellular retinol-binding protein form.

[provided by RefSeq, Jul 2008]

Synonyms: ALDH1A6; ALDH6; MCOP8; RALDH3






Protein Families: Druggable Genome

Protein Pathways: Drug metabolism - cytochrome P450, Glycolysis / Gluconeogenesis, Histidine metabolism,


Metabolic pathways, Metabolism of xenobiotics by cytochrome P450, Phenylalanine

metabolism, Tyrosine metabolism

Product images:





HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY ALDH1A3 (Cat# [RC209656], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ALDH1A3 (Cat# [TA502841])(1:500).



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-ALDH1A3 monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human).







Negative control E. coli lysate (Left lane) or E. coli lysate containing recombinant protein fragment for human ALDH1A3 (NP_000648) gene (the fusion of amino acids 1-100 and 413-512) (Right lane). Equivalent amounts (5 ug per lane) were separated by SDS-PAGE and then immunoblotted with anti-ALDH1A3.

Flow cytometric Analysis of Jurkat cells, using anti-ALDH1A3 antibody ([TA502841]), (Red), compared to a nonspecific negative control antibody, (Blue).