

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA502849BM

PNMT Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI1E8]

Product data:

| Product Type: | Primary Antibodies |
|-------------------------|---|
| Clone Name: | OTI1E8 |
| Applications: | FC, IF, WB |
| Recommended Dilution: | WB 1:2000, IF 1:100, FLOW 1:100 |
| Reactivity: | Human, Mouse |
| Host: | Mouse |
| lsotype: | lgG2b |
| Clonality: | Monoclonal |
| Immunogen: | Full length human recombinant protein of human PNMT (NP_002677) produced in HEK293T cell. |
| Formulation: | PBS (pH 7.3) containing 1% BSA, 50% glycerol. |
| Concentration: | 0.5 mg/ml |
| Purification: | Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G) |
| Conjugation: | HRP |
| Storage: | Store at -20°C as received. |
| Stability: | Stable for 12 months from date of receipt. |
| Predicted Protein Size: | 30.7 kDa |
| Gene Name: | phenylethanolamine N-methyltransferase |
| Database Link: | <u>NP_002677</u> <u>Entrez Gene 18948 MouseEntrez Gene 5409 Human</u> <u>P11086</u> |
| Background: | The product of this gene catalyzes the last step of the catecholamine biosynthesis pathway, which methylates norepinephrine to form epinephrine (adrenaline). The enzyme also has beta-carboline 2N-methyltransferase activity. This gene is thought to play a key step in regulating epinephrine production. [provided by RefSeq] |
| Synonyms: | PENT; PNMTase |




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


PNMT Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI1E8] – TA502849BM

Protein Families: Druggable Genome

Protein Pathways: Metabolic pathways, Tyrosine metabolism

Product images:






HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PNMT ([RC206586], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PNMT. Positive lysates [LY400946] (100ug) and [LC400946] (20ug) can be purchased separately from OriGene.

Anti-PNMT mouse monoclonal antibody ([TA502849]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY PNMT ([RC206586]).

HEK293T cells transfected with either [RC206586] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-PNMT antibody ([TA502849]), and then analyzed by flow cytometry.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Flow cytometric Analysis of Jurkat cells, using anti-PNMT antibody ([TA502849]), (Red), compared to a nonspecific negative control antibody, (Blue).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US