

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA502917S

AP2M1 Mouse Monoclonal Antibody [Clone ID: OTI1C12]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI1C12
Applications:	FC, WB
Recommended Dilution:	WB 1:2000, FLOW 1:100
Reactivity:	Human, Mouse, Rat
Host:	Mouse
lsotype:	lgG2a
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human AP2M1 (NP_001020376) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	0.63 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	49.2 kDa
Gene Name:	adaptor related protein complex 2 subunit mu 1
Database Link:	<u>NP_001020376</u> <u>Entrez Gene 11773 MouseEntrez Gene 116563 RatEntrez Gene 1173 Human Q96CW1</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

GRIGENE AP2M1 Mouse Monoclonal Antibody [Clone ID: OTI1C12] – TA502917S


Background:	This gene encodes a subunit of the heterotetrameric coat assembly protein complex 2 (AP2),
	which belongs to the adaptor complexes medium subunits family. The encoded protein is
	required for the activity of a vacuolar ATPase, which is responsible for proton pumping
	occurring in the acidification of endosomes and lysosomes. The encoded protein may also
	play an important role in regulating the intracellular trafficking and function of CTLA-4
	protein. Two transcript variants encoding different isoforms have been found for this gene.
	[provided by RefSeq]

Synonyms:	AP50; CLAPM1; mu2
Protein Families:	Druggable Genome
Protein Pathways:	Endocytosis, Huntington's disease

Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY AP2M1 ([RC201377], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-AP2M1. Positive lysates [LY422484] (100ug) and [LC422484] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-AP2M1 monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human) (1:200).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



HEK293T cells transfected with either [RC201377] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-AP2M1 antibody ([TA502917]), and then analyzed by flow cytometry.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US