

## Product datasheet for **TA502930AM**

### **S6K1 (RPS6KB1) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI4G4]**

#### **Product data:**

Product Type:	Primary Antibodies
Clone Name:	OTI4G4
Applications:	FC, IF, WB
Recommended Dilution:	WB 1:2000, IF 1:100, FLOW 1:100
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human RPS6KB1 (NP_003152) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Biotin
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	59 kDa
Gene Name:	ribosomal protein S6 kinase B1
Database Link:	<a href="#">NP_003152</a> <a href="#">Entrez Gene 72508 Mouse</a> <a href="#">Entrez Gene 83840 Rat</a> <a href="#">Entrez Gene 6198 Human</a> <a href="#">P23443</a>



[View online »](#)

**Background:**

This gene encodes a member of the RSK (ribosomal S6 kinase) family of serine/threonine kinases. This kinase contains 2 non-identical kinase catalytic domains and phosphorylates several residues of the S6 ribosomal protein. The kinase activity of this protein leads to an increase in protein synthesis and cell proliferation. Amplification of the region of DNA encoding this gene and overexpression of this kinase are seen in some breast cancer cell lines. Alternate translational start sites have been described and alternate transcriptional splice variants have been observed but have not been thoroughly characterized. [provided by RefSeq]

**Synonyms:**

p70 S6KA; p70(S6K)-alpha; p70-alpha; p70-S6K; PS6K; S6K; S6K-beta-1; S6K1; STK14A


**Protein Families:**

Druggable Genome, Protein Kinase


**Protein Pathways:**

Acute myeloid leukemia, ErbB signaling pathway, Fc gamma R-mediated phagocytosis, Insulin signaling pathway, mTOR signaling pathway, TGF-beta signaling pathway


**Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY RPS6KB1 ([RC217324], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-RPS6KB1. Positive lysates [LY401097] (100ug) and [LC401097] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (10ug) from 2 different cell lines by using anti-RPS6KB1 monoclonal antibody (1:200).



Anti-RPS6KB1 mouse monoclonal antibody ([TA502930]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY RPS6KB1 ([RC217324]).



HEK293T cells transfected with either [RC217324] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-RPS6KB1 antibody ([TA502930]), and then analyzed by flow cytometry.