

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA502960AM

EB2 (MAPRE2) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI3C7]

Product data:

Product Type:	Primary Antibodies	
Clone Name:	OTI3C7	
Applications:	FC, IF, WB	
Recommended Dilution:	WB 1:500~2000, IF 1:100, FLOW 1:100	
Reactivity:	Human, Dog, Rat, Monkey	
Host:	Mouse	
lsotype:	lgG2b	
Clonality:	Monoclonal	
Immunogen:	Full length human recombinant protein of human MAPRE2 (NP_055083) produced in HEK293T cell.	
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.	
Concentration:	0.5 mg/ml	
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)	
Conjugation:	Biotin	
Storage:	Store at -20°C as received.	
Stability:	Stable for 12 months from date of receipt.	
Predicted Protein Size:	36.9 kDa	
Gene Name:	microtubule associated protein RP/EB family member 2	
Database Link:	<u>NP 055083</u> <u>Entrez Gene 679221 RatEntrez Gene 490487 Dog</u> Entrez Gene 707996 MonkeyEntrez Gene <u>10982 Human</u> <u>Q15555</u>	



	EB2 (MAPRE2) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI3C7] – TA502960AM
--	--


Background:

The protein encoded by this gene shares significant homology to the adenomatous polyposis coli (APC) protein-binding EB1 gene family. The function of this protein is unknown; however, its homology suggests involvement in tumorigenesis of colorectal cancers and proliferative control of normal cells. This gene may belong to the intermediate/early gene family, involved in the signal transduction cascade downstream of the TCR. Alternative splicing results in multiple transcript variants. [provided by RefSeq]


Synonyms:	EB1; EB2; RP1
Protein Families:	Druggable Genome

Protein Families:


Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY MAPRE2 ([RC200259], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-MAPRE2. Positive lysates [LY415395] (100ug) and [LC415395] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-MAPRE2 monoclonal antibody.




Anti-MAPRE2 mouse monoclonal antibody ([TA502960]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY MAPRE2 ([RC200259]).



HEK293T cells transfected with either [RC200259] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-MAPRE2 antibody ([TA502960]), and then analyzed by flow cytometry.



Flow cytometric Analysis of Hela cells, using anti-MAPRE2 antibody ([TA502960]), (Red), compared to a nonspecific negative control antibody (TA50011), (Blue).



Flow cytometric Analysis of Jurkat cells, using anti-MAPRE2 antibody ([TA502960]), (Red), compared to a nonspecific negative control antibody (TA50011), (Blue).