

Product datasheet for TA502963

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

SNAP25 Mouse Monoclonal Antibody [Clone ID: OTI4G5]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI4G5

Applications: FC, IF, IHC, WB

Recommended Dilution: WB 1:2000, IHC 1:150, IF 1:100, FLOW 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human SNAP25 (NP_003072) produced in

HEK293T cell.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.9 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 23.2 kDa

Gene Name: synaptosome associated protein 25

Database Link: NP 003072

Entrez Gene 20614 MouseEntrez Gene 25012 RatEntrez Gene 6616 Human

P60880





Background:


Synaptic vesicle membrane docking and fusion is mediated by SNAREs (soluble Nethylmaleimide-sensitive factor attachment protein receptors) located on the vesicle membrane (v-SNAREs) and the target membrane (t-SNAREs). The assembled v-SNARE/t-SNARE complex consists of a bundle of four helices, one of which is supplied by v-SNARE and the other three by t-SNARE. For t-SNAREs on the plasma membrane, the protein syntaxin supplies one helix and the protein encoded by this gene contributes the other two. Therefore, this gene product is a presynaptic plasma membrane protein involved in the regulation of neurotransmitter release. Two alternative transcript variants encoding different protein isoforms have been described for this gene. [provided by RefSeq]

Synonyms: bA416N4.2; dJ1068F16.2; RIC-4; RIC4; SEC9; SNAP; SNAP-25


Protein Families: Druggable Genome

Protein Pathways: SNARE interactions in vesicular transport

Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY SNAP25 (Cat# [RC202068], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-SNAP25 (Cat# TA502963). Positive lysates [LY418912] (100ug) and [LC418912] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffinembedded Human colon tissue within the normal limits using anti-SNAP25 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.






Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human colon tissue using anti-SNAP25 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.




Immunohistochemical staining of paraffinembedded Human liver tissue within the normal limits using anti-SNAP25 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.




Immunohistochemical staining of paraffinembedded Carcinoma of Human lung tissue using anti-SNAP25 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.






Immunohistochemical staining of paraffinembedded Carcinoma of Human pancreas tissue using anti-SNAP25 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.




Immunohistochemical staining of paraffinembedded Carcinoma of Human bladder tissue using anti-SNAP25 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Anti-SNAP25 mouse monoclonal antibody (TA502963) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY SNAP25 ([RC202068]).





HEK293T cells transfected with either [RC202068] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-SNAP25 antibody (TA502963), and then analyzed by flow cytometry.